

TECHNICAL DATA SHEET FOLIMIX (CITRIC ACID CHELATED MICRONUTRIENT MIX)

DESCRIPTION: Water Soluble Citric Acid Chelated Micro Element Mixture Fertilizer composed of **Iron, Zinc, Manganese, Copper and Boron for agricultural use only.**

TECHNICAL SPECIFICATIONS:

Physical Form: Water Soluble Powder Fertilizer
Product Colour: Light Green
Product Odour: Odourless
Shelf Life: 4 years
Origin: TURKEY
Producer: SAMİ KÜÇÜKKOCA TARIM TİCARET

COMPOSITION (w/w):

Ferrous (Fe) 6.0 %
Zinc (Zn) 5.0 %
Manganese (Mn) 3.0 %
Copper (Cu) 3.0 %
Boron (B) 1.0 %
Ph: 2 - 4

ADVANTAGES AND PROPERTIES :

FOLIMIX is recommended for application to all crops and plants.
FOLIMIX supplies Iron, Zinc, Manganese, Boron, Copper demand of plants.
FOLIMIX improves quality and productivity. Solve plant's basic Micronutrient problems.
FOLIMIX is a non – phytotoxic micronutrient for both foliar and soil application.

DIRECTIONS FOR USE :

-Apply 1 Kg. to 2 Kg. per Hectare for foliar or 2 Kg. to 4 Kg. per Hectare for soil application within 3 weeks after emergence. Split applications are preferred over a single application. Timing and rates for split applications are 1 to 1.5 Kg. foliar or 2 to 3 Kg. banded within 3 weeks after emergence, and repeat application before bloom.

FRUIT AND NUT CROPS: Almonds, Apples and Pears, Apricots, Nectarines, Peaches, Plums and Prunes, Cherries, Citrus, Grapes, Pecans...

SOIL ... 3 - 10 Kg./Hectare, with pesticide, irrigation or fertilizer.

FOLIAGE... Mix 2 Kg. to 3 Kg. per Hectare with sufficient water and apply as a through cover spray. Repeat At petal fall and cover sprays at 1 Kg. to 2 Kg. per Hectare. Do not apply more than 2 Kg. per Hectare to plums or sensitive varieties with cover sprays. For Citrus, Mix 3 Kg. to 5 Kg. per Hectare with sufficient water and apply during spring and fall growth flush. Repeat during growing Season at 1 Kg. to 2 Kg. per Hectare, with pesticide cover sprays.

FIELD AND VEGETABLE CROPS: Beans, Corn, Cotton, Potatoes, Rice, Wheat, Barley, Oats, Sorghum...

SOIL & FOLIAGE ... Apply 1 Kg. to 2 Kg. per Hectare to the foliage or soil. For best results, split application as follows. Apply 1 Kg. to 1.5 Kg. per Hectare to the foliage or soil (band) within 3 weeks of emergence. Apply 0.5 Kg. to 1 Kg. per Hectare with each pesticide application prior to bloom.

PACKING : 0.25 - 0.5 - 1 - 2 - 5 - 10 - 25 Kg.

APPROPRIATE WAREHOUSE CONDITIONS: Try to keep the product between + 5, - 25 C in its original packaging far from the human, animal and food. Be aware of direct sunlight.

USAGE SECURITY REGULATIONS : This product is not toxic. After usage wash your hands and face with water and soap.